
Brandon J. Erickson, MD Mackenzie Lindeman, ATC 176 3rd Ave New York, NY 658 White Plains Rd Tarrytown, NY 450 Mamaroneck Rd Harrison, NY

Phone: 914-580-9624

Brandon.erickson@rothmanortho.com Mackenzie.lindeman@rothmanortho.com

https://rothmanortho.com/physicians/brandon-j-erickson-md



Snapping Scapula Debridement Physical Therapy Protocol

NameDate
Diagnosis s/p RIGHT/LEFT Debridement of Snapping Scapula
Date of Surgery
Frequency:times/week Duration: Weeks
Weeks 1-4:
Sling for four weeks
Pendulums allowed after first postoperative visit Encourage hand, wrist, elbow ROM, grip strengthening
Encourage nand, wrist, cloow KOW, grip strengthening
Weeks 4-6:
Begin AAROM, isometrics, for rotator cuff, deltoid
ADLs around the house without sling
Continue to wear sling in public for 6 weeks
Washa 6.2 Marshar
Weeks 6-3 Months: Global AROM
Theraband Exercise for rotator cuff, deltoid, scapular rotators
Expectation: Return to work and sport at full activity averaged 3.5 to 4.5 months
Comments:
Functional Capacity EvaluationWork Hardening/Work Conditioning Teach HEF
Modalities
Electric StimulationUltrasound IontophoresisPhonophoresisTENS
Heat beforeIce afterTrigger points massage Therapist's discretion
Signature